研究業績 英文表記

表題 The relationship between children's somatotypes, motor examination results, and motor skills: assessing 6 to 10-year-olds 著者名 久保温子、村田伸、安彦鉄平、田中真一 所属 西九州大学リハビリテーション学部 正 英文 Title The relationship between children's somatotypes, motor examination results, and motor skills: assessing 6 to 10-year-olds Author Atsuko Kubo, Shin Murata, , Teppei Abiko,Shinichi Tanaka, Affiliation Faculty of Rehabilitation Science, Nishikyushu University Author Atsuko Kubo, Shin Murata, , Teppei Abiko,Shinichi Tanaka, Affiliation Faculty of Rehabilitation Science, Nishikyushu University Abstract IPurposel Childhood motor disorders and obesity are major health problems and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of for motor examination items, Physical Fitness Test, and body composition analysis for two groups. [Results] More obese children were unable to reach the floor while performing side-torside jumping, and obese children showed better values. Many endomorphic children were also unable to touch the floor while performing side-torside jumping, and obese children showed better values. Many endomorphic children were also unable to tou	和文	
所属 西九州大学リハビリテーション学部 度文 Title The relationship between children's somatotypes, motor examination results, and motor skills: assessing 6 to 10-year-olds Author Atsuko Kubo., Shin Murata, , Teppei Abiko,Shinichi Tanaka, Affiliation Faculty of Rehabilitation Science, Nishikyushu University Purposel Childhood motor disorders and obesity are major health problems and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of four motor examination items, Physical Fitness Test, and body composition analysis for two groups. [Results] More obese children were unable to reach the floor while performing a standing forward bend compared to non obese children. In addition, a significant difference was found in the assessment of motor performance while performing isde-to-side jumping, and obese children showed better values. Many endomorphic children were also unable to touch the floor while performing side to-side jumping, and bese children showed better values. Many endomorphic children were also unable to touch the floor while performing side to-side jumping, and obese children showed better values. Many endomorphic children were also unable to touch the floor while performing side to-side jumping, and obese children showed better values. Many endomorphic children were also unable to touch the floor while performing side to-side jumping, and obese children shomed better values	表題	
技文 Title The relationship between children's somatotypes, motor examination results, and motor skills: assessing 6 to 10-year-olds Author Atsuko Kubo,, Shin Murata, , Teppei Abiko, Shinichi Tanaka, Affiliation Faculty of Rehabilitation Science, Nishikyushu University Image: Participants and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of four motor examination in terms. Physical Fitness Test, and body composition analysis for two groups. Results! More obese children were unable to reach the floor while performing a standing forward bend compared to non-obese children. In addition, a significant difference was found in the assessment of motor performance while performing the to side to side pos revealed significant differences. There were no somatotype related differences in the results of the body composition analysis. [Conclusion] In children aged significant differences was found in the assessment of motor performance while performing the standing forward bend. Among the floor with their hands when performing the standing forward bend. Among the floor performance while performing the standing forward bend. Among the floor with their hands when performing the standing forward bend. Among the floor shill perform the state to side to side hops percealed significant differences. There were no somatotype related differences in the results of the body composition analysis. [Conclusion] In children aged six to ten yea	著者名	久保温子、村田伸、安彦鉄平、田中真一
Title The relationship between children's somatotypes, motor examination results, and motor skills' assessing 6 to 10-year-olds Author Atsuko Kubo, Shin Murata, , Teppei Abiko, Shinichi Tanaka, Affiliation Faculty of Rehabilitation Science, Nishikyushu University Image: Purposel Childhood motor disorders and obesity are major health problems and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of four motor examination items. Physical Fitness Test, and body composition analysis for two groups. [Results] More obese children were unable to reach the floor while performing a standing forward bend compared to non-obese children. In addition, a significant differences was found in the assessment of motor performance while performing the standing forward bend compared to touch the floor with their hands when performing the standing forward bend. Among the items from a physical fitness test, the side-to-side hops revealed significant differences. There were no somatotype related differences in the results of the body composition analysis. [Conclusion] In children aged six to ten years, somatotype differences were not associated with motor skill or body composition.	所属	西九州大学リハビリテーション学部
Author Atsuko Kubo,, Shin Murata, , Teppei Abiko, Shinichi Tanaka, Affiliation Faculty of Rehabilitation Science, Nishikyushu University Image: Purposel Childhood motor disorders and obesity are major health problems and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of four motor examination items, Physical Fitness Test, and body composition analysis for two groups. [Results] More obese children were unable to reach the floor while performing a standing forward bend compared to non-obese children. In addition, a significant difference was found in the assessment of motor performance while performing the standing forward bend. Among the items from a physical fitness test, the side-to-side hops revealed significant differences. There were no somatotype related differences in the results of the body composition analysis. [Conclusion] In children aged six to ten years, somatotype differences were not associated with motor skill or body composition.	英文	
Affiliation Faculty of Rehabilitation Science, Nishikyushu University IPurpose] Childhood motor disorders and obesity are major health problems and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of four motor examination items, Physical Fitness Test, and body composition analysis for two groups. [Results] More obese children were unable to reach the floor while performing a standing forward bend compared to non-obese children. In addition, a significant difference was found in the assessment of motor performance while performing the standing forward bend. Among the items from a physical fitness test, the side to side hops revealed significant differences. There were no somatotype-related differences in the results of the body composition analysis. [Conclusion] In children aged six to ten years, somatotype differences were not associated with motor skill or body composition.	Title	
Abstract Abstra	Author	Atsuko Kubo,, Shin Murata, , Teppei Abiko,Shinichi Tanaka,
Abstract and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of four motor examination items, Physical Fitness Test, and body composition analysis for two groups. [Results] More obese children were unable to reach the floor while performing a standing forward bend compared to non-obese children. In addition, a significant difference was found in the assessment of motor performance while performing side-to-side jumping, and obese children showed better values. Many endomorphic children were also unable to touch the floor with their hands when performing the standing forward bend. Among the items from a physical fitness test, the side-to-side hops revealed significant differences. There were no somatotype-related differences in the results of the body composition analysis. [Conclusion] In children aged six to ten years, somatotype differences were not associated with motor skill or body composition.	Affiliation	Faculty of Rehabilitation Science, Nishikyushu University
keyword Children, Somatotype, Motor examinatio	Abstract	and concerns in children today. We performed a physical examination to test the motor system and motor ability of elementary school children based on their body types. [Participants and Methods] The obesity levels of 161 elementary school students aged six to ten were calculated based on the gender, age, and standard weight for each height category to classify them into somatotype groups, and analyze the relationships among the results of four motor examination items, Physical Fitness Test, and body composition analysis for two groups. [Results] More obese children were unable to reach the floor while performing a standing forward bend compared to non-obese children. In addition, a significant difference was found in the assessment of motor performance while performing side-to-side jumping, and obese children showed better values. Many endomorphic children were also unable to touch the floor with their hands when performing the standing forward bend. Among the items from a physical fitness test, the side-to-side hops revealed significant differences. There were no somatotype-related differences in the results of the body composition analysis. [Conclusion] In children aged six to ten years, somatotype differences were not associated with motor skill or body
	keyword	Children, Somatotype, Motor examinatio

※本データの英文表記は実際の論文上の表記とは異なります。