Ittle Community Author Yuki Kishikawa, et al. Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences Image: The validity of a quality of life (QOL) scale was examined in elderesidents of a community. Twenty-eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture w included, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v answered "not healthy" in Ota QOL were significantly lower th those of those who answered "healthy" in QOL. The part correlation ratio, which indicates the relative contribution of factores			
著者名 岸川曲紀、他 所属 西九州大学リハビリテーション学部リハビリテーション学科 英文 Title Validity of the Quality of Life Assessment Scale for the Elderly Living in Community Author Yuki Kishikawa, et al. Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences Abstract The validity of a quality of life (QOL) scale was examined in elderesidents of a community. Twenty-eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture wincluded, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v answered "not healthy" in QoL and WHO QOL. The part correlation ratio, which indicates the relative contribution of fact was '0.62. These results suggest that the Ota QOL scale has valid	和文		
所属 西九州大学リハビリテーション学部リハビリテーション学科 英文 Title Validity of the Quality of Life Assessment Scale for the Elderly Living in Community Author Yuki Kishikawa, et al. Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences Abstract The validity of a quality of life (QOL) scale was examined in elderesidents of a community. Twenty eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture w included, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v answered "not healthy" in Ota QOL were significantly lower the those of those who answered "healthy" in QOL. The part correlation ratio, which indicates the relative contribution of fact was -0.62. These results suggest that the Ota QOL scale has valid	表題	地域在住高齢者に対する QOL 評価尺度の妥当性	
英文 Title Validity of the Quality of Life Assessment Scale for the Elderly Living in Community Author Yuki Kishikawa, et al. Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences Image: State of the stat	著者名	岸川由紀、他	
Title Validity of the Quality of Life Assessment Scale for the Elderly Living in Community Author Yuki Kishikawa, et al. Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences The validity of a quality of life (QOL) scale was examined in elderesidents of a community. Twenty-eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture w included, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v answered "not healthy" in Ota QOL were significantly lower th those of those who answered "healthy" in QOL. The par correlation ratio, which indicates the relative contribution of factor was -0.62. These results suggest that the Ota QOL scale has valid	所属	西九州大学リハビリテーション学部リハビリテーション学科	
Ittle Community Author Yuki Kishikawa, et al. Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences The validity of a quality of life (QOL) scale was examined in elderesidents of a community. Twenty-eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture wincluded, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v answered "not healthy" in Ota QOL were significantly lower th those of those who answered "healthy" in QOL. The parcorrelation ratio, which indicates the relative contribution of factor was -0.62. These results suggest that the Ota QOL scale has valid	英文		
Affiliation Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences The validity of a quality of life (QOL) scale was examined in elderesidents of a community. Twenty-eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture w included, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v answered "not healthy" in Ota QOL were significantly lower the those of those who answered "healthy" in QOL. The part correlation ratio, which indicates the relative contribution of factor was -0.62. These results suggest that the Ota QOL scale has valid	Title	Validity of the Quality of Life Assessment Scale for the Elderly Living in the Community	
AbstractThe validity of a quality of life (QOL) scale was examined in elderesidents of a community. Twenty-eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture wincluded, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v answered "not healthy" in Ota QOL were significantly lower the those of those who answered "healthy" in QOL. The part correlation ratio, which indicates the relative contribution of factor was -0.62. These results suggest that the Ota QOL scale has valid	Author	Yuki Kishikawa, et al.	
Abstractresidents of a community. Twenty-eight males and females aged years or older (mean age 78.7 years) living in Saga Prefecture w included, and the SF-8, Ota QOL, and WHO QOL26 were used QOL scales. The results showed that the scores of those v 	Affiliation	Department of Rehabilitation Sciences, Faculty of Rehabilitation Sciences	
	Abstract	The validity of a quality of life (QOL) scale was examined in elderly residents of a community. Twenty-eight males and females aged 65 years or older (mean age 78.7 years) living in Saga Prefecture were included, and the SF-8, Ota QOL, and WHO QOL26 were used as QOL scales. The results showed that the scores of those who answered "not healthy" in Ota QOL were significantly lower than those of those who answered "healthy" in QOL. The partial correlation ratio, which indicates the relative contribution of factors, was -0.62. These results suggest that the Ota QOL scale has validity as a QOL scale for the elderly.	
keyword Quality of life	keyword	Quality of life	

研究業績 英文表記

※本データの英文表記は実際の論文上の表記とは異なります。