## 研究業績 英文表記

和文	
表題	リハビリテーション利用者における手段的日常生活動作と主観的健康の関係
著者名	宮原洋八 ¹, Makoto Uchinoura², 岸川由紀 ¹, Yuji Yamaguchi², 小松洋平 ¹, 松尾萌美 ¹, Seijiro Nishimura², Shinichi Shibata², Hajime Shibata²
所属	「西九州大学リハビリテーション学部, <sup>2</sup> 久留米リハビリテーション病院
英文	
Title	Relationship between instrumental activities of daily living and self-rated health rehabilitation users
Author	Hiroya Miyabara <sup>1</sup> , Makoto Uchinoura <sup>2</sup> , Yuki Kishikawa <sup>1</sup> , Yuji Yamaguchi <sup>2</sup> , Youhei Komatsu <sup>1</sup> , Moemi Matsuo <sup>1</sup> , Seijiro Nishimura <sup>2</sup> , Shinichi Shibata <sup>2</sup> , Hajime Shibata <sup>2</sup>
Affiliation	<sup>1</sup> Faculty of Rehabilitation Sciences, Nishi Kyushu University, <sup>2</sup> Kurume Rehabilitation Hospital
Abstract	In general, elderly persons with limitations in instrumental activities of daily living (IADL) are fragile, due to their various health problems such as disorders, poor cognitive functions. Physical and cognitive status could be predictors for the onset of IADL limitation in elderly persons. On the other hand, self-rated health (SRH), a valid generic health measurement is used as a resource and as an outcome, also was reported as a reliable measurement because it was verified that results of SRH and mortality were correlated. Thus, it can be deemed as a resource which affecting inclusive and dynamic health behavior, and reflecting measure of health status. Therefore, our main objective was to clarify relationships between limitation of IADL levels, SRH, and Mini-Mental State Examination (MMSE) among elderly persons who use ambulatory rehabilitation. As the result, IADL limitation was related to the level of SRH, but not with MMSE scores.
keyword	Instrumental activities of daily living, Self-rated health, Rehabilitation users, The long-term care insurance