

## 研究業績 英文表記

和文	
表題	地域在住高齢者の自己申告による杖の使用に関連する要因の検討
著者名	釜崎大志郎 <sup>1,2</sup> , 大田尾浩 <sup>1</sup> , 八谷瑞紀 <sup>1</sup> , 久保温子 <sup>1</sup> , 大川裕行 <sup>1</sup> , 坂本飛鳥 <sup>1</sup> , 藤原和彦 <sup>1</sup> , 保坂公大 <sup>3</sup> , 北島貴大 <sup>4</sup> , 下木原俊 <sup>2,5</sup> , 丸田道雄 <sup>2,6</sup> , 韓伋熙 <sup>2,7</sup> , 溝上泰弘 <sup>8</sup> , 鎌田實 <sup>9</sup> , 田平隆行 <sup>10</sup>
所属	<ol style="list-style-type: none"> <li>1. 西九州大学リハビリテーション学部</li> <li>2. 鹿児島大学医学部客員研究員</li> <li>3. 久留米リハビリテーション病院 リハビリテーション科</li> <li>4. ひらまつ病院リハビリテーション科</li> <li>5. 札幌医科大学保健医療学部作業療法学科</li> <li>6. 長崎大学大学院医歯薬学総合研究科作業療法学専攻</li> <li>7. 国際医療福祉大学福岡医療保健学部作業療法学科</li> <li>8. 株式会社 MIZ</li> <li>9. 諏訪中央病院</li> <li>10. 鹿児島大学大学院保健学研究科</li> </ol>
英文	
Title	Examination of factors associated with self-reported cane use among community-dwelling older adults
Author	Taishiro Kamasaki <sup>1,2</sup> Hiroshi Otao <sup>1</sup> Mizuki Hachiya <sup>1</sup> Atsuko Kubo <sup>1</sup> Hiroyuki Okawa <sup>1</sup> Asuka Sakamoto <sup>1</sup> Kazuhiko Fujiwara <sup>1</sup> Kodai Hosaka <sup>3</sup> Takahiro Kitajima <sup>4</sup> Suguru Shimokihara <sup>2,5</sup> Michio Maruta <sup>2,6</sup> Gwanghee Han <sup>2,7</sup> Yasuhiro Mizokami <sup>8</sup> Minoru Kamata <sup>9</sup> Takayuki Tabira <sup>10</sup>
Affiliation	<ol style="list-style-type: none"> <li>1. Department of Rehabilitation Sciences, Nishikyushu University</li> <li>2. Visiting Researcher, Faculty of Medicine, Kagoshima University</li> <li>3. Medical Corporation Kabutoyama-kai Kurume Rehabilitation Hospital Rehabilitation Center</li> <li>4. Medical Corporation Hiramatsu Hospital</li> <li>5. Department of Occupational Therapy, School of Health Sciences, Sapporo Medical University</li> <li>6. Department of Occupational Therapy, Nagasaki University Graduate School of Biomedical Sciences</li> <li>7. Department of Occupational Therapy, School of Health Sciences at Fukuoka, International University of Health and Welfare</li> <li>8. Miz Co., Ltd</li> <li>9. Suwa Central Hospital</li> <li>10. Graduate School of Health Sciences, Kagoshima University</li> </ol>
Abstract	This study investigates factors influencing cane use among older adults and identifies most significant determinants. There were 108 participants (77 ± 7 years) in the cane-non-using group and 52 (83 ± 6 years) in the cane-using group. Factors associated with the presence or absence of cane use were open-eyed one-leg standing time [OR 0.81 (0.70–0.93), p = 0.003] and depressed mood [OR 2.78 (1.31–5.91), p = 0.008]. Older adults with reduced balance ability and depressive mood need to use a cane. This highlights the need to assess balance ability and depressive mood in prescribing a cane to older adults. Appropriate prescription of canes has the potential to enrich the lives of older adults by contributing to their safe range of activities.
keyword	Walking aids; Open-eyed one-leg stand time; Depressive moods; Older adults

※本データの英文表記は実際の論文上の表記とは異なります。